

Effect of Grain Geometry on Angle of Repose and Dynamics

C. J. Olson, C. Reichhardt, M. McCloskey, and R. J. Zieve
Department of Physics, University of California, Davis, California 95616.
(February 1, 2008)

With simulation and experiment we examine the effect of grain anisotropy on the angle of repose in a two-dimensional geometry. We find that the structure of the granular packing can be directly controlled by changing the aspect ratio of the grains, and that the specific type of order strongly affects the angle of repose and the resulting dynamics. The angle of repose is largest for dimer grains, and is smaller for both monomer and trimer grains. The motion of a collapsing grain heap consists of large scale collective shear when monomer grains are considered, but occurs via tumbling of grains on the surface for dimers, and via disorderly bulk tumbling for trimer grains.

PACS numbers: 45.70.Mg, 81.05.Rm, 45.70.-n, 45.50.-j

Granular assemblies have attracted much recent attention since they are a model system in which collective nonequilibrium dynamics, compaction, pattern formation, frustration, nonlinear flows, and jamming can be readily examined [1]. The behavior they display is relevant to a diverse array of physical systems, including powders, colloids, liquids, and vortex matter in superconductors. In addition, an understanding of granular mechanics will be valuable for increasing the efficiency of a wide range of industrial processes and applications. A fundamental question that arises is the relation between the mechanical response of the granular media and the ordering of the individual grains. The nature of the close-packed ordering in 3D of different types of granular media is itself an open issue. In this work we study a simple system of granular assemblies confined in a 2D plane and show that specific types of ordering can be controllably produced by tuning the anisotropy of the grains. The resulting structure of the granular pack has a profound effect on the angle of repose and the granular dynamics.

Most previous work on the effects of grain geometry has focused on angles of repose for grains that are roughly spherical. The effect of dispersion in the sizes of individual spherical grains has been considered [2], as well as the effect of varying the roughness of the grain surface [3] or of having nonspherical grains [4,5]. Anisotropic or elongated grains have attracted interest previously, but it has generally been in the context of the flow of feed grains through a hopper [6]. Only one systematic study has considered the effect of gradual elongation of the grains on the angles of repose. In Ref. [5], it was observed that as the grain is elongated continuously from a monomer to a dimer, the angle of repose increases monotonically. The grain shape was continuously varied, so it would not be possible to form an ordered lattice with the same lattice constant using any two grain geometries considered.

In this work we consider with experiments and simulations a system composed of grains that can always be packed into a triangular lattice of monomers, but are individually composed of monomers, dimers, or trimers. Some previous simulation studies have focused on fully three-dimensional systems [7,8]. This introduces consid-

erable complexity to the problem since there are many more degrees of freedom, and it is difficult to determine experimentally the behavior of the grains below the surface of the pile [9]. For this reason, we consider a system constrained to move in two dimensions. Here the microscopic behavior of individual grains can be directly observed experimentally and compared with computer simulations.

With experiments and simulations we find that the maximum angle of stability depends on the grain aspect ratio in a non-monotonic way. Both monomers, which are fully ordered, and trimers, which are both positionally and bond orientationally disordered, are less stable than dimers, which have positional order but not bond-orientational order. We directly link these specific types of ordering to the grain dynamics, which range from orderly large scale shear flow for monomers, to tumbling surface flow for dimers, to disordered bulk flow for trimers. Our results in 2D represent a first step towards the general understanding of how the microscopic structure affects the mechanical response in 3D granular assemblies.

Experiment—We consider a system of steel balls of radius $1/16''$ constrained to move in 2D by two Plexiglas sheets [10]. The steel balls are either used individually, or are welded together to form dimers or linear trimers (see inset to Fig. 1). All three shapes can tile space in triangular arrays with the same lattice constant. The grains are poured into a vertical enclosed area 9" wide by 14" high, and the container is briefly shaken vertically to prepare a more ordered initial starting condition. Monomers and dimers form nearly perfect triangular lattices under these conditions, but there is always a large amount of disorder present in the trimers [10]. The container is slowly tilted in the plane of the Plexiglas sheets to change the angle between the surface of the granular packing and gravity [2]. We record the angle at which the first grain motion occurs, giving us the maximum angle of stability Θ_s for the system, which is similar to the angle of repose.

In Fig. 1 we show the appearance of a system of dimers after an avalanche has occurred. The grains remained motionless until the container had been tipped by 60° ,




FIG. 1. Digital camera image of experimental appearance of a system of dimers after an avalanche has occurred. Patches of order remain present in the system. Inset: Experimental monomers, dimers, and trimers.

and then a large avalanche occurred. A significant amount of order remains in the system even after the avalanche, as seen in Fig. 1. The maximum angles of stability Θ_s measured experimentally in this way for grains of differing anisotropy are listed in Table I. We find the lowest angle $\Theta_s = 43^\circ$ for the disordered trimer system. The well-ordered monomers have a slightly higher value of $\Theta_s = 45^\circ$, which is in reasonable agreement with previous experiments on spherical or nearly spherical grains [2,3]. In contrast dimers display a much larger angle of stability of $\Theta_s = 62^\circ$. To better understand the reasons that the dimer shape is so stable, we turn to a simulation of the system.

Simulation—We consider a system of $N_g = 705$ grains constrained to move in a 2D plane in a system of size 10×18 to 60×18 . We use a granular dynamics method similar to the one employed in [11–13] to integrate the equations of motion for each grain, given by: $m_i \mathbf{v}_i = \mathbf{f}_{\text{el}}^{(i)} + \mathbf{f}_{\text{diss}}^{(i)} + \mathbf{f}_{\text{shear}}^{(i)} + \mathbf{f}_g + \mathbf{f}_{\text{fric}}$. Here, $\mathbf{f}_g = -0.0025$ is the force of gravity and $\mathbf{f}_{\text{fric}} = 0.3$ is the friction between the grain and the plane in which the grains move (representing the Plexiglas plate). Two grains interact only when their relative distance is smaller than the sum of their radii, $r_g = 0.4$. In this regime, three forces are active. The elastic restoration force between two grains is $\mathbf{f}_{\text{el}}^{(i)} = \sum_{i \neq j} k_g m_i (|\mathbf{r}_{ij}| - 2r_g) \mathbf{r}_{ij} / |\mathbf{r}_{ij}|$, where $k_g = 20$ is the strength of the restoring spring, $2r_g$ is the grain diameter, $m_i = 1$ is the grain mass and $\mathbf{r}_{ij} = \mathbf{r}_j - \mathbf{r}_i$, the distance between grains located at $\mathbf{r}_i$ and $\mathbf{r}_j$. The dissipation force due to the inelasticity of the collision is $\mathbf{f}_{\text{diss}}^{(i)} = -\sum_{i \neq j} \gamma m_i (\mathbf{v}_{ij} \cdot \mathbf{r}_{ij}) \mathbf{r}_{ij} / |\mathbf{r}_{ij}|^2$, where $\gamma = 2.4$ is a phenomenological dissipation coefficient and $\mathbf{v}_{ij} = \mathbf{v}_i - \mathbf{v}_j$ is the relative velocity. The shear friction force mimicking solid friction is $\mathbf{f}_{\text{shear}} = -\gamma_s m_i (\mathbf{v}_{ij} \cdot \mathbf{t}_{ij}) \mathbf{t}_{ij} / |\mathbf{r}_{ij}|^2$, where $\gamma_s = 1.2$ is the shear friction coefficient and $\mathbf{t}_{ij} = (-r_{ij}^y, r_{ij}^x)$ is the vector $\mathbf{r}_{ij}$ rotated by 90° . The behavior we observe is not sensitive to

our choice of parameters.

In this model, we neglect Coulomb friction and the rotation of individual particles. In some cases, we constrain the grains to form dimers or trimers by rigidly fixing the grains together. These composite dimers and trimers are free to rotate. The walls and floor are evenly lined with immobile grains of the same size as the simulated monomers. Such a rough floor is necessary to produce a finite angle of repose in the case of monomers; without it, the monomers continue to slump until their height is reduced to a single grain. Dimers and trimers, however, produce a finite angle of repose even on a smooth floor due to their tendency to interlock.

The system is prepared in one of two ways. In studies of Θ_s , the grains are prepared in a box of size 10×18 such that the top surface of the grains is roughly flat, either by dropping individual grains from above, or by placing them into an ordered arrangement directly. In studies of the dynamics of collapsing grains, the grains are first introduced into a box that is $1/6$ as wide as the full simulation area of 60×18 , either by dropping or by direct placement. The right wall is then instantaneously moved out to the edge of the system, and the grains collapse to the right. This method has been previously employed in Refs. [14,15] to study angles of repose.

When the grains are dropped slowly from above, we find that monomers and dimers fall into well-ordered arrangements but trimers are disordered. This is illustrated in Fig. 2, where the positions of the grains are shown in Fig. 2(a,c,e), and the Delaunay triangulations of the lattice are shown in Fig. 2(b,d,f). The monomers of Fig. 2(a,b) are perfectly ordered, with no dislocations, as indicated in Table I. For dimers as in Fig. 2(c,d) the positional order is nearly complete but there is a slight chance for a vacancy to occur. Note that although the individual elements composing the dimers are ordered, the bond angles have no overall order. For trimers, shown in Fig. 2(e,f), there is some local ordering of the bond angles, but no overall order, and there is a large amount of positional disorder in the lattice.

There is a simple geometric reason for the difference in packing of the three grain shapes. If a layer of previously dropped grains is present, and there is a single monomer-sized vacancy in the layer, then an additional monomer or dimer dropped over the vacancy will fall into the vacancy and fill it. The dimer is not long enough to span the vacancy and one side of it will slip in. In contrast, the trimer is long enough to span the vacancy and block it from being filled. Additionally, a trimer is long enough to assume a continuum of angles if dropped near a second trimer lying flat, depending on the exact relative spacing of the two trimers, whereas a dimer will assume only the angles allowed by a triangular lattice. Thus with trimers only very local ordering is possible and long-range order cannot be achieved through random dropping.

We next consider Θ_s , measured with the same




FIG. 2. Static images from simulation after grains have been dropped into a container. Monomers: (a) positions, (b) Delaunay triangulation. The system is well ordered. Dimers: (c) positions, with lines connecting the dimer elements, (d) Delaunay triangulation, with dislocation sites colored grey. The positions of the individual grains are ordered, but there is no order in the orientation of the dimer bonds. Trimers: (e) positions, with lines connecting the trimer elements, (f) Delaunay triangulation. There is disorder in both position and bond orientation.

procedure followed in the experiment. We simulate tipping the box slowly by rotating the direction of gravity, and record the angle Θ_s at which the first avalanche occurs. As listed in Table I, we find the same trend as experiment: low values of $\Theta_s = 27^\circ$ for ordered monomers and $\Theta_s = 13^\circ$ for disordered trimers, and a much larger angle of $\Theta_s = 40^\circ$ for the dimers. It is clear that the different types of order in these three geometries lead to the different angles of stability. The disordered trimers are very unstable to rearrangements since there are many dislocations in the granular packing where motion can occur. The ordered monomers are more stable due to




FIG. 3. Dynamic images of granular collapse from simulation at pairs of consecutive time intervals after the right wall of the container has been removed. Circles indicate the location of individual grain elements, light lines indicate bonds for dimers and trimers, and heavier lines indicate the trajectories over a short time of the individual grain elements. (a,b) Monomers, showing orderly flow along lattice vector directions of large triangular wedges. (c,d) Dimers, showing tumbling flow along the top surface of the pile only. (e,f) Trimers, showing disorderly flow throughout the bulk, with tumbling motion at the top of the pile.

the absence of dislocations, but are still unstable to shearing motion along the lattice directions. In contrast, the dimers are stabilized by the absence of dislocations like the monomers, but shearing motion is suppressed due to the randomly arranged bonds between the dimers, which lead to interlocking. The larger Θ_s we observe for dimers compared to monomers agrees with the results of Ref. [5]. Our study of the even more anisotropic trimers indicates, however, that the increase in Θ_s with anisotropy found in Ref. [5] does not continue indefinitely but instead decreases again as the grains become longer than dimers.

To further explore how the structure of the granular packing affects its stability, we directly examine the dynamics of collapsing grains when one wall is removed, and compare the three geometries in Fig. 3. We see a striking difference in the dynamics. Collapsing monomers move in large collective motions involving much of the bead pack, which follow the lattice vectors. The motion is very orderly and occurs in the form of large pulses, during which triangular wedges of grains displace along their




FIG. 4. Net grain velocity in the x direction, V_x , from simulation. Top (heavy) line: Monomers, showing large collective oscillations. Middle line: Disordered dimers. Bottom (light) line: Trimers. The curve for the monomers has been vertically offset by 0.4 and the curve for dimers has been offset by 0.2 for clarity.

lattice vectors. This wedge-like motion is similar to motion seen in experiments with flowing steel balls [16].

In collapsing dimers, the motion is limited to the surface of the pile due to the interlocking of the dimers which prevents shearing motion along the lattice vectors. The motion that does occur is tumbling in nature and very disorderly. A relatively small portion of the bead pack is moving at any given time.

In collapsing trimers, the large number of dislocations present in the system serve as “lubrication” to the motion, allowing easy disorderly sliding of a large portion of the bead pack. There is no significant ordering present on a scale of more than a few grains so coherent motion is impossible. Motion occurs through a combination of gliding along the dislocated area, and tumbling down the topmost portion of the pile. Although the motion is widespread, it occurs more slowly than the collective motion seen in the monomers. This is due to the fact that a considerable amount of rearrangement within the bulk must occur as the movement continues, unlike the case of monomers when no rearrangement was necessary and the movement could follow the lattice vectors.

The different dynamics can also be observed by measuring the net velocity in the x -direction, $V_x = \sum v_x$, of the grains over time, as shown in Fig. 4. Monomers produce extremely large pulses as the collective motion occurs, whereas dimers and trimers produce much smaller amplitude signals with their more disperse responses.

Summary— We have examined the ordering and dynamics of monomer, dimer and trimer assemblies of grains confined in 2D and relate the specific types of ordering to the dynamics of granular collapse. Dimers, which have translational order but not bond-orientational order, show the largest angle of repose.

Monomers, which are completely ordered, show a smaller angle of repose that is still higher than trimers which are completely disordered. The dynamics of the monomers during collapse occurs by large scale collective shear motions. The dimers, which do not allow for easy shear motion show a surface tumbling motion. Trimers also show a tumbling motion that can occur in the bulk as well as on the surface.

Acknowledgements: We thank H. Jaeger and N. Grønbech-Jensen for helpful discussions. This work was supported in part by CLC and CULAR (LANL/UC) and by NSF, DMR-9733898.

- [1] H. Jaeger, S.R. Nagel, and R.P. Behringer, *Rev. Mod. Phys.* **68**, 1259 (1996).
- [2] F. Cantelaube *et al.*, *J. Phys. I (France)* **5**, 581 (1995).
- [3] C.M. Dury *et al.*, *Phys. Rev. E* **57**, 4491 (1998); J.P. Koeppe, M.ENZ, and J. Kakalios, *ibid.* **58**, R4104 (1998); H.M. Jaeger, C. Liu, and S.R. Nagel, *Phys. Rev. Lett.* **62**, 40 (1989).
- [4] J.A. Cuesta and D. Frenkel, *Phys. Rev. A* **42**, 2126 (1990); B.J. Buchalter and R.M. Bradley, *ibid.* **46**, 3046 (1992); T. Poschel and V. Buchholtz, *Phys. Rev. Lett.* **71**, 3963 (1993); D. Hirshfeld, Y. Radzyner, and D.C. Rapaport, *Phys. Rev. E* **56**, 4404 (1997); F.X. Villarruel *et al.*, *ibid.* **61**, 6914 (2000).
- [5] J.A.C. Gallas and S. Sokolowski, *Int. J. Mod. Phys. B* **7**, 2037 (1993).
- [6] G.W. Baxter and R.P. Behringer, *Phys. Rev. A* **42**, 1017 (1990).
- [7] J. Baxter, U. Tuzuen, J. Burnell, and D.M. Heyes, *Phys. Rev. E* **55**, 3546 (1997).
- [8] Y.C. Zhou, B.D. Wright, R.Y. Yang, B.H. Xu, and A.B. Yu, *Physica A* **269**, 536 (1999).
- [9] T.G. Drake, *J. Geophys. Res.* **95**, 8681 (1990).
- [10] I.C. Rankenburg and R. Zieve, preprint.
- [11] J.A.C. Gallas, H.J. Herrmann, and S. Sokolowski, *Phys. Rev. Lett.* **69**, 1371 (1992).
- [12] H.J. Herrmann, *Int. J. Mod. Phys. C* **4**, 309 (1993).
- [13] S. Luding, E. Clement, A. Blumen, J. Rajchenbach, and J. Duran, *Phys. Rev. E* **50**, 4113 (1994).
- [14] T. Elperin and E. Golshtein, *Physica A* **242**, 332 (1997).
- [15] J. Lee and H.J. Herrmann, *J. Phys. A* **26**, 373 (1993).
- [16] T.G. Owe Berg, R.L. McDonsals, and R.J. Trainor, *Powder Technol.* **3**, 56 (1969).

Grain type	Θ_s (Expt.)	Θ_s (Sim.)	P_d (Expt.)	P_d (Sim.)
Monomer	45°	27°	0%	0%
Dimer	62°	40°	3%	3%
Trimer	43°	16°	22%	30%

TABLE I. Table of results from experiment and simulation. The angle of stability Θ_s was obtained by tipping the container and identifying the first motion of the flat granular surface. P_d is the percentage of individual grain elements that do not have six neighbors (0 indicates perfect order).